

General Surgery at Northpointe

| Gail Strindberg, MD | | Sarah Johnson, NP-C |

2376 N 400 E Ste 205 Tooele, Utah 84074

Phone: (435)882-8111

Northpointe Surgical Center

2326 N 400 E Bldg. C Ste 100

Tooele, Utah 84074

(435) 843-0180

Mountain West Medical Center

2055 N Main St

Tooele, Utah 84074

(435) 843-3600

PATIENT INSTRUCTIONS FOR COLONOSCOPY

Procedure Date: _____

PHYSICIAN: Gail Strindberg, MD

_____ NO SOLID FOOD Yes ___ NO ___ Blood draw 90 days prior

**** If you need to reschedule or cancel, please contact our office as soon as possible.**

THE FACILITY WILL CALL YOU THE AFTERNOON BEFORE YOUR SCHEDULED PROCEDURE WITH YOUR SCHEDULED TIME.

**** A couple days before procedure, drink more water each day, at least 8 cups ****

Day before procedure:

Drink only clear liquids for breakfast, lunch, and dinner. **NO SOLID FOODS.**

In the morning mix the container of Golytely/Trilyte by adding a gallon of water or Gatorade (**no red or purple**) and place in the refrigerator to chill.


In the evening, begin your prep at 6:00 pm. Drink **HALF** of the container (2 liters) by drinking 8 oz. every 10 minutes. Rapid drinking of the solution is better than drinking small amounts continually. Continue to drink **clear liquids** until bedtime, **HYDRATION** is extremely important for a successful colonoscopy.

Morning of the procedure:

Drink the 2nd half (2 liters) of the Golytely 5-6 hours before procedure (example: if your procedure time is 9 am, start drinking the Golytely at 3-4 am). Every 10 minutes drink 8 oz of solution until the container of Golytely is empty. DO NOT drink any liquids 3 hours before your procedure.

- You will receive an IV sedation and **MUST have someone drive you home.**
- We recommend, flushable wipes and Vaseline in place of toilet paper, to avoid becoming too sore.
- YOU MAY STAY ON YOUR ASPIRIN. Ask about other blood thinners such as Plavix/Clopidogrel. Some of them may need to be stopped 5 days before your procedure.

Clear Liquids Allowed:



Anything you can see through, such as water, clear fruits juices (apple and white grape) any flavor bouillon, Jell-O, clear soft drinks, Gatorade or Powerade

Clear Liquids you MAY have the DAY BEFORE your test:

- Water
- Ice
- Coffee or Tea (no creamer or milk, sugar is ok)
- Soft Drinks (Orange, Ginger Ale, Coke, Sprite & 7UP) (**NOTHING red or purple**)
- Sports Drinks, Kool-Aid, Crystal Light (Orange or Lemonade, (**NOTHING red or purple**))
- Strained Fruit Juice, **PULP FREE** (Apple Juice, White Grape, White Cranberry & Lemonade.
- Low Salt Chicken or Beef Broth (Clear Broth or Bouillon)
- Hard Candies (**NOTHING red or purple**)
- Jell-o (Lemon, Lime & Orange, (**NOTHING red or purple**))
- Popsicles and Italian Ice, (**NOTHING red or purple**, no ice cream, sherbet, yogurt or fruit bars)

Items NOT allowed the DAY BEFORE your test:

- **NO** Alcoholic Beverages
- **NOTHING** Red or Purple
- **NO** dairy
- **NO** Solid Foods

*******NOTHING THE DAY OF PROCEDURE*******

Please note:

If you need to take any medications the morning of the procedure, it is recommended to take them with **ONLY 2 Tablespoons or less of WATER.**

Any Questions, please call our office (435)882-8111

Insurance Coverage for Colonoscopy

A screening colonoscopy is for patients 45 years or older with no current or previous issues. This should be covered 100% by your insurance.

If you have hemorrhoids removed with your screening, you may owe those charges.

If you have polyps, personal or family history of polyps/cancer, it is no longer considered a screening, but a diagnostic procedure and you may be charged for your colonoscopy.

If you owe for any of your procedure, you may receive a billing statement from:

1. General Surgery at Northpointe (Dr. Strindberg)
2. Northpointe Surgical Center
3. Deseret Peak Anesthesia
4. Diagnostic Laboratory